

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/12, 39/13, 39/15, 39/145, 39/29, C12N 5/06, 7/02		A1	(11) International Publication Number: WO 96/16672 (43) International Publication Date: 6 June 1996 (06.06.96)
(21) International Application Number: PCT/US95/15118 (22) International Filing Date: 30 November 1995 (30.11.95) (30) Priority Data: 08/351,079 30 November 1994 (30.11.94) US (71) Applicants: DYNACORP [US/US]; Suite 200, One Taft Court, Rockville, MD 20850 (US). NATIONAL INSTITUTES OF HEALTH [US/US]; 9000 Rockville Pike, Bethesda, MD 20892 (US). (72) Inventors: CHANOCK, Robert, M.; 7001 Longwood, Bethesda, MD 20817 (US). PURCELL, Robert, H.; 17517 White Ground Road, Boyds, MD 20720 (US). KAPIKIAN, Albert, Z.; 11201 Marcliff Road, Rockville, MD 20852 (US). POTASH, Louis; 5337 Pooks Hill Road, Bethesda, MD 20814 (US). (74) Agents: STEPNO, Norman, H. et al.; Burns, Doane, Swecker & Mathis, P.O. Box 1404, Alexandria, VA 22313-1404 (US).		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>	
(54) Title: SERIALY PASSAGED AFRICAN GREEN MONKEY KIDNEY CELLS			
(57) Abstract The invention described herein provides a method for selecting a novel African Green monkey kidney (AGMK) cell substrate, its cultivation and serial passage and its subsequent characterization. The invention further provides a method for the use of the cell substrate in the isolation, growth and serial passage of a large number of viruses, particularly rotaviruses, enteroviruses, respiratory viruses and hepatitis A virus. A method for the utilization of the AGMK cell substrate for the production of live and killed virus vaccines is also provided.			
<div style="text-align: right;"> Ref. #2 SLU 4538 R. Belshe et al. 09/785,764 </div>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

SERIALLY PASSAGED AFRICAN GREEN MONKEY KIDNEY CELLS

**STATEMENT AS TO RIGHTS TO THE INVENTION MADE UNDER
FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT**

Part of the work performed during development of this invention utilized U.S. Government funds. The U.S. Government has certain rights in this invention (National Institute of Allergy and Infections Diseases Contract No. NO1-AO-35154).

FIELD OF INVENTION

The invention relates to a novel African Green Monkey Kidney (AGMK) cell substrate that supports the efficient replication of human and animal rotaviruses, astroviruses, enteroviruses including the Sabin live attenuated poliovirus vaccine strains, hepatitis A virus and respiratory viruses such as respiratory syncytial virus, parainfluenza viruses, and influenza A and B viruses.

The invention is also useful for the production of virus suspensions suitable for use in vaccines for immunization of humans.

15 BACKGROUND OF INVENTION

Rotaviruses are major causal agents of acute gastroenteritis in man with a world-wide distribution. Enteroviruses have been implicated in infections of the gastrointestinal tract, the respiratory system and in aseptic meningitis. Respiratory syncytial virus (RSV), the parainfluenza viruses and the influenza viruses have been shown to be the major causal agents of serious pediatric diseases such as croup, bronchiolitis and pneumonia, while the influenza viruses

are the major cause of serious febrile respiratory tract illness in adults. Hepatitis A virus is the cause of hepatitis A, one of the major infectious diseases of the liver.

Rotavirus infections occur world-wide and are responsible for a large proportion of severe, life-threatening, often fatal diarrheal disease. Most rotavirus disease is caused by the four major human rotavirus serotypes but other serotypes are being discovered continuously. The latter are usually restricted in geographic distribution, but they could become a larger problem at any time. World-wide, the human rotaviruses are the major etiologic agents of serious acute gastroenteritis in infants and young children and on occasion can cause debilitating diarrhea in adults. It has been estimated by the World Health Organization that rotaviruses are responsible for one million fatal diarrheal illnesses each year in infants and young children in developing countries.

The most important cause of serious viral respiratory illness in children is respiratory syncytial virus (RSV). The parainfluenza viruses, influenza viruses, and adenoviruses are also important in this regard. Especially in the case of respiratory syncytial virus (RSV), of which there are two major subgroups A and B, immunity does not appear to be long-lasting and re-infections can and do occur with high frequency during infancy, through the pre-school period, and throughout adult life. The parainfluenza viruses, of which there are four major types -- 1, 2, 3 and 4 -- have been implicated as important causes of croup, bronchiolitis and pneumonia in infants and young children. These viruses are

second only to RSV as a cause of severe viral respiratory tract disease. In adults, the influenza viruses are a major cause of mortality in older persons with underlying acute or chronic cardiac or pulmonary disease. The serious systemic disease manifestations of influenza virus infections are well known and the frequent antigenic shifts in serotype [A/H1N1, A/H3N2 and B] necessitate changes in the composition of the strains included in the yearly vaccine preparations.

The enteroviruses, which are a subgroup of picomaviruses consisting of polioviruses, Coxsackie viruses and echoviruses, have been shown to cause a broad spectrum of illnesses. These illnesses include paralytic disease, encephalitis, aseptic meningitis, pleurodynia, exanthems, and pericarditis with some of the infections resulting in debilitating sequelae. Hepatitis A virus, also a picomavirus, is a major cause of sporadic as well as epidemic hepatitis.

According to convention, semi-continuous cell systems which are diploid and have a finite longevity in terms of the number of passages that can be achieved in the laboratory are designated as cell strains, whereas continuous cell systems that are aneuploid and can be passaged indefinitely in the laboratory (i.e., are immortal) are designated as cell lines. A limiting factor in the development of vaccines for combating the maladies caused by a number of these infectious agents has been the lack of availability of an acceptable cell strain or line capable of supporting the growth of these viruses to a concentration suitable for use in vaccine production. Convenient and susceptible cell strains and/or lines that can be cultured using relatively high split ratios (i.e., the preparation of

multiple tissue culture vessels from a single culture) and which can be approved for human use by the Center for Biologics Evaluation and Research (CBER) of the Food and Drug Administration (FDA) are needed to enable and facilitate vaccine development for certain medically important viruses that are uncontrolled
5 currently.

Semi-continuous or continuous cell culture systems capable of supporting the growth of hybrid rotaviruses, hereafter designated human x animal rotavirus reassortants, have included the simian cell strain, FRhL-2, and the simian cell lines, CV-1 and Vero. Live virus vaccines produced in these cells have received
10 CBER, FDA approval for Phase I and II studies. Henceforth, we will refer to these cell strains or lines as certified cell strains or lines to distinguish them from cell strains or lines that have been licensed for production of human vaccines. To date, only two semi-continuous cell culture systems, WI-38 and MRC-5, both human fetal diploid cell strains, have been licensed for use in virus vaccines.
15 These cell strains will be referred to as licensed cell strains. Currently, there are no licensed continuous cell lines.

The above mentioned certified cell strain or lines (FRhL-2, CV-1 and Vero) are limited in their ability to support the growth of completely homologous human rotaviruses, i.e., rotaviruses that derive each of their 11 RNA gene
20 segments from a human rotavirus. In addition, the FRhL-2 cell strain, with a maximum 1:3 split ratio capability, is limited in its ability to support the growth of important RSV. The CV-1 cells, with a maximum 1:4 split ratio capability, have not supported the growth of many viral agents to a level satisfactory for

vaccine production. The Vero cells, with a split ratio of 1:6 - 1:10, are limited in their ability to support the growth of a number of enteroviruses and fail to support the efficient growth of completely homologous human rotaviruses.

Naturally occurring strains of hepatitis A virus (HAV), a picomavirus
5 distantly related to the enteroviruses, do not grow well in any cell type during
primary isolation and must be adapted to growth in cells of primate origin before
the level of virus replication required for vaccine development and production
can be achieved. Inactivated whole-virus HAV vaccines grown in fetal human
fibroblast MRC-5 cells have been developed, but the low split ratio of 1:2
10 required by these cells, the low viral titers achieved, and the prolonged
cultivation time of up to two weeks necessary for maximum yield of viral antigen
make such vaccines expensive. Growth of HAV in simian CV-1, FRhK-4 (a
fetal rhesus monkey kidney cell line), FRhK-6 (another fetal rhesus monkey
kidney cell line) and Vero cells has been variable depending on the strain and
15 passage level of virus, but generally is suboptimal. Best growth has been
obtained in a cloned cell line, designated clone 11-1, derived from FRhK-4 cells,
but these cells contain bovine papilloma virus sequences and thus, are not
suitable for vaccine development.

Candidate live attenuated and inactivated HAV vaccines have been
20 developed by adaptation of the virus to growth in primary African green monkey
(AGMK) cells but such vaccines are not economically feasible because of the
extreme difficulty of obtaining sufficient primary AGMK cells that are free of
extraneous viral agents of monkey origin. Live attenuated and inactivated HAV

vaccine candidates have been adapted to growth in MRC-5 cells, a human fetal diploid cell strain, but such adaptation has consistently led, respectively, to over-attenuation of the live attenuated virus for humans and a relatively sparse yield of inactivated virus vaccine.

5 Thus, there remains a need for a cell strain or line capable of supporting the efficient growth of a large number of human viral pathogens such as the human rotaviruses, enteroviruses, HAV, and respiratory viruses of major medical importance including RSV, influenza viruses and parainfluenza viruses. Such a cell strain or line would facilitate the development and production of
10 commercially useful and effective vaccines. There is a pressing need for the development of vaccines against these viruses in order to prevent severe viral diseases of infants, children, adults and the elderly.

 Therefore, the availability of a single cell strain or line capable of supporting the efficient replication of those viral agents responsible for a
15 significant number of childhood and adult diseases would provide a significant advancement in the formulation and production of multivalent vaccines.

SUMMARY OF THE INVENTION

 The present invention is directed to a cell substrate that substantially overcomes the limitations of previously established cell strains or lines. The
20 serially passaged cells of the present invention are derived from the paired kidneys of an African green monkey and support the efficient growth of medically important human rotaviruses, astroviruses, picomaviruses such as

enteroviruses including the Sabin live attenuated poliovirus vaccine strains and hepatitis A virus, and respiratory viruses such as respiratory syncytial virus, parainfluenza viruses, and influenza A and B viruses.

The AGMK cells of the invention are designated as a cell substrate
5 because it is not clear at this time whether they constitute a cell strain or cell line. It is probable that they are a cell line because they are aneuploid, an essential and defining property of cell lines. However, the AGMK cells have been passaged serially only 30 times and assignment of cell immortality, one of the other defining features of a cell line, requires more than 50 serial passages.

10 At the 30th passage there was no diminution of viability, or ability to grow nor was there any other outward sign of senescence. Furthermore, the AGMK cells have been split 1:4 or 1:8 at each of these passages, a split ratio that is characteristic of cell lines but not cell strains which usually can not be split more than 1:2.

15 An important advantage of the present invention is that the cell substrate, based on currently promulgated guidelines (i.e., "Points to Consider in the Characterization of Cell Lines Used to Produce Biologicals" (May 1993)) and state-of-the-art testing, has been demonstrated to be free of detectable adventitious microbial agents. This feature allows the cell substrate to be
20 employed for the growth of viruses, such as the human rotaviruses and hepatitis A virus, which have a very highly restricted and narrow tissue culture host range. The unavailability of such a clean cell system prior to this invention has hampered vaccine development for these viruses. In addition, the cell substrate

can be used in studies which require cultures of monkey kidney cells free of simian adventitious agents. The commercially available monkey kidney cell cultures are frequently contaminated with these agents, which negate or becloud any results obtained and preclude their use in vaccines destined for administration to humans. This invention also offers the advantage of a cell substrate which can be trypsinized and split in ratios ranging from 1:4 to 1:8 depending upon vessel size and time constraints imposed by vaccine production deadlines.

Additional features and advantages of the invention will be set forth in the description which follows or may be learned by practice of the invention. The objectives and other advantages of the invention to be realized and attained will be cited in the written description and claims hereof as well as the appended drawings.

To achieve these and other advantages and in accordance with the purpose of the invention, as embodied and broadly described, the invention provides a cell line derived from the paired kidneys of an African green monkey, which is free of demonstrable adventitious microbial agents. The invention also provides a method for establishing the cell line by the enzymatic disaggregation of the paired kidneys from an African green monkey and continuous cultivation of the resulting cells. In addition, the invention provides a method for recovering and serially propagating specific viruses in the cell line that subsequently can be used for production of a viral vaccine.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a cell line derived from a pair of African green monkey kidneys. The cell line is free of demonstrable viable adventitious microbial agents. In addition, the cell line is capable of supporting the growth of
5 viruses such as human rotaviruses, astroviruses, enteroviruses, respiratory viruses and hepatitis A and is, therefore, useful for propagating viruses necessary for the production and formulation of a number of effective vaccines on a scale that makes such vaccines commercially feasible.

Specifically, the present invention relates to an aneuploid cell substrate
10 with chromosome counts in the diploid range which was initiated by the enzymatic disaggregation of paired kidneys from a female *Cercopithecus* African green monkey. A deposit of the cell line at passage No. 13, Lot 6500 has been made with the American Type Culture Collection, 12301 Parklawn Drive, Rockville, MD 20852, USA, on November 4th and has been assigned accession
15 No. ATCC# CRL 11756.

The present invention also provides a method of establishing a cell substrate capable of at least 30 serial passages that is initially derived by the enzymatic disaggregation of kidneys from an African green monkey. In addition, the invention provides a method for recovery and growth of specific viruses, a
20 necessary first step for subsequent vaccine production.

It will be apparent to those skilled in the art that various modifications and variations can be made in the cell substrate of the present invention and in its uses without departing from the spirit or scope of the invention. Thus, it is

intended that the present invention covers the modifications and variations of this invention provided they come within the scope of the appended claims and their equivalents.

Having now described the invention in general terms, it will be better
5 understood by reference to certain examples which are included herein for the purpose of illustration only and are not intended to be limiting unless otherwise specified.

EXAMPLE 1

Derivation of the AGMK Cell Substrate

10 Source: The cell substrate was derived from the kidneys of an African green monkey from a colony that had been established on Barbados. The cells are free of any demonstrable viable adventitious microbial agents. The microbial agents include, but are not limited to bacteria, fungi, mycobacteria, mycoplasma and simian agents such as retroviruses, foamy viruses, hemadsorbing viruses, herpes-
15 like viruses and multi-nucleating or syncytial viruses. The kidneys were processed using conventional enzymatic disaggregation technology. The resulting kidney cell suspension was used to initiate 3 primary flask [T150 cm²] cultures with the remainder, designated as Freeze #2129, distributed into multiple ampules and frozen for storage in a liquid nitrogen freezer. The primary flask
20 cultures labelled A, B and C, were processed as follows:

Flask A: On day 14 split 1:2 = 1-A-1 & 1-A-2 with one of these
flasks split on day 28 1:2 = 2-A-1 & 2-A-2

Flask B: On day 21 split 1:2 = 1-B-1 & 1-B-2 with one of these
flasks split on day 35 1:2 = 2-B-1 & 2-B-2

Flask C: Not split

The maintenance medium consisted of Eagle's Minimal Essential Medium
5 (EMEM) + 5% fetal bovine serum (heat inactivated at 56°C for 30 min.) + 1%
glutamine (200 mM). All flask cultures were refed with 100 ml of fresh
maintenance medium on a weekly basis for a total incubation period of 9 weeks
based on the initial planting date. The cultures were negative for hemadsorption
and CPE (cytopathic effect), the latter determined by microscopic examination
10 after fixation and staining.

The passage history from a single ampule to the preparation of the Master
Seed Cell Bank (MSCB) and the Master Working Cell Bank (MWCB) is depicted
in Figure 1. Cultures were handled in a Class 100 Laminar Flow Hood and in
an area where only normal tissue cultures and/or vaccine-designated cell systems
15 were cultured. Cultures were serially passaged employing a 1:4 to 1:6 split ratio
with incubation at 36°C ± 1°C and utilizing the following methodology:

1. Cell films were washed with Hanks' Balanced Salt Solution
(HBSS) without Ca⁺⁺ and Mg⁺⁺.
2. Films were disaggregated by the addition of a trypsin-EDTA
20 solution and incubation at 36°C ± 1°C for 4-5 minutes.
3. Films were broken up by rapid pipetting with the cells suspended
in the growth medium [EMEM + 10% fetal bovine serum + 1%
glutamine] and seeded into appropriate size culture vessels.

The passage history from single ampules from the MWCB to the post-production testing is outlined in Figure 2.

EXAMPLE 2

Testing of the Master Working Cell Bank - Passage 7A, Lot #4214

5

1. **KARYOTYPING AND SPECIES IDENTIFICATION:** Passage 7A, Lot 94214 was characterized by Giemsa banded chromosome analysis and reaction with species-specific antisera and isoenzyme analysis.

a. **Cytogenic Characterization:** Conventional chromosome staining (non-banded) was utilized to determine chromosome count ploidy distribution per 100 metaphases. Identifiable markers and aberrations were analyzed for 50 metaphases.

Chromosome banding by trypsin-Giemsa was utilized to analyze the karyotype and identify specific chromosome markers. Species confirmation was made using G-banding patterns and chromosome morphology. The procedures used were modifications of those described by Peterson, W.D., Jr., Simpson, W.F., and Hukku, B., Cell culture characteristics: Monitoring for cell identification, in Jakoby, W.B. and Pastan, I.H., eds., *Methods in Enzymology* 58:164-178, 1979; and Seabright, M., A rapid banding technique for human chromosomes, *Lancet*, ii:971-972, 1971.

20

The cell line was identified as an aneuploid female *Cercopithecus* (African green) monkey (XX) with chromosome counts in the diploid range. Most of the

chromosomes were normal and present in two copies each. Normal chromosome D was monosomic and a single marker D chromosome was observed.

b. **Species Identification of Cell Culture:** The ability of the test cell preparation to react with species-specific antisera was determined using the

5 fluorescent antibody technique as described by Peterson *et al.*, *supra*. The cells reacted with monkey antiserum but not with mouse or hamster antiserum.

c. **Isoenzyme Analysis:** Enzyme preparations were made from the test article cells. The electrophoretic migration distances of the enzymes were compared with known migration distances of enzymes in various mammalian

10 species as described by Halton, D.M., Peterson, W.D., Jr.; and Hukku, B. Cell culture quality control by rapid isoenzymatic characterization, *In Vitro* 19:16-24, 1983; Ottenbreit, M.J., Halton, D.M., and Peterson, W.D., Jr., Rapid isoenzyme analysis of cell cultures by agarose electrophoresis, I. Interspecies identification, *J. Tissue Culture Methods*, 6:107-110, 1982; and II. Intraspecies
15 identification of human cell lines, *J. Tissue Culture Methods*, 8:125-130, 1986.

The electrophoretic mobilities of enzymes glucose-6-phosphate dehydrogenase (G6PD), nucleoside phosphorylase (NP), malate dehydrogenase (MDH) and lactate dehydrogenase (LDH) present in an extract prepared from the test cells were comparable to those of a *Cercopithecus* (African green) monkey
20 control cell preparation. No extra bands were observed on the electrophoresis films that might suggest the presence of cells of a species other than that of the cell culture.

Cells other than those of the submitted culture (MWCB, Passage 7A, Lot# 4214) were not detected in the culture by either isoenzyme analysis or by cytogenic evaluation.

2. DETECTION OF RETROVIRUSES

5 a. Detection of Retrovirus Particles by Electron Microscopic

Examination: A cell pellet of the test article cells (MWCB, Passage 7A, Lot# 4214) was prepared and examined for the presence of virus particles by transmission electron microscopy. No retrovirus-like particles were found in the 200 cells observed.

10 b. Detection of Retrovirus Reverse Transcriptase in the Presence of

DNA-Dependent DNA Polymerase: The test article (MWCB, Passage 7A, Lot# 4214) was analyzed for the presence of types B, C and D retrovirus reverse transcriptase and cellular DNA polymerase activity. No evidence of the presence of retrovirus reverse transcriptase was observed.

15 The results are presented in Table I - A, B and C.

TABLE I-A

Reverse Transcriptase and DNA Polymerase Activities
in Test Article AGMK MWCBC Cell Cultures: p7A, L-4214

Sample	Adjusted Reverse Transcriptase ^a (rAdT)	Reverse Transcriptase ^b (rAdT)		DNA Polymerase ^b (dAdT)	
	Mn ⁺⁺	Mn ⁺⁺	Mn ⁺⁺	Mn ⁺⁺	Mn ⁺⁺
<i>Test Article</i>					
Undiluted	22	25	0	7	0
Diluted 2-fold in medium ^c	6	8	0	19	1
Diluted 2-fold in R-MuLV ^d		20,837	433	10	0
Diluted 2-fold in DNA polymerase ^e		795	6	7,137	508
<i>Positive and Negative Controls^f</i>					
R-MuLV diluted 2-fold in medium ^g		18,549	549	12	0
DNA Polymerase diluted 2-fold in medium ^g		1,079	25	5,443	542
SMRV ^h		2,585	48,289	211	396
Stabilization Buffer ⁱ		(63)	(59)	(67)	(58)
Medium ^c		(65)	(57)	(65)	(52)
Medium ^g		(78)	(56)	(62)	(49)

^a Reverse transcriptase activity adjusted for the effect of DNA polymerase (Table 2 and Methods).

^b Expressed as counts per minute ³H-thymidine triphosphate incorporated (mean duplicates minus background).

^c EMEM + 10% serum subtracted from test article samples.

^d R-MuLV (Rauscher Murine Leukemia Virus) - Type C virus positive control.

^e DNA polymerase - Positive control; prepared from mink lung cells.

^f Control monitoring background of incorporation of ³H-thymidine triphosphate (reaction mixture with undiluted test article minus template primers) = 61 cpm (medium background not subtracted).

^g EMEM + 10% serum subtracted from R-MuLV and DNA polymerase positive controls.

^h SMRV (Squirrel Monkey Retrovirus Preparation) - Type D virus positive control.

ⁱ Stabilization buffer used to dilute SMRV positive control.

TABLE I-B

Adjustment of Reverse Transcriptase Values for Interference by Polymerase
in Test Article AGMK MWCB Cell Cultures: p7A, L-4214

	RT Assay (CPM)		DNA Polymerase Assay (CPM)	
	Observed ^a (Mn ⁺⁺)	Adjusted ^b (Mn ⁺⁺)	Observed (Mn ⁺⁺)	DNA Polymerase x Slope
<i>Test Article</i>				
Undiluted	25	22	37	3
Diluted 2-fold in medium ^c	8	6	19	2

^a Unadjusted ³H-TTP incorporation catalyzed by RT. May include effects by cellular DNA polymerase.

^b Adjusted ³H-TTP incorporation catalyzed by RT = Observed RT - (DNA polymerase x Slope). Slope = 0.08.

^c Medium used to dilute test article.

TABLE I-C

Reverse Transcriptase and DNA Polymerase Activities
in a DNA Polymerase Preparation from Reverse
Transcriptase-Negative Cell Cultures

(Used for Determining Interference in Reverse
Transcriptase Assay by DNA Polymerase)

DNA Polymerase ^a	Reverse Transcriptase ^b (rAdT)		DNA Polymerase ^b (dAdT)	
	Mn ⁺⁺	Mg ⁺⁺	Mn ⁺⁺	Mg ⁺⁺
Sample 1	116	10	1,898	198
Sample 2	187	0	2,634	0
Sample 3	547	6	6,690	119
Sample 4	574	3	8,529	81
Sample 5	1,196	27	14,315	1,222

^a DNA polymerase prepared from mink lung cells.

^b Expressed as counts per minute ³H-thymidine triphosphate incorporated [mean duplicates minus background (medium)].

EXAMPLE 3**Post-Production Cell Testing****1. Karyotyping and Species Identification:**

The AGMK cells at Passage 17A, Lot #4706 were characterized by
5 Giemsa banded chromosome analysis and reaction with species-specific antisera
and isoenzyme analysis.

The cells were identified as aneuploid female *Cercopithecus* (African
green) monkey (XX, XXX) with chromosome counts in the hypodiploid to
hyperdiploid range. All karyotypes examined contained a single normal
10 *Cercopithecus* monkey marker chromosome (Group D) and a second partially
deleted *Cercopithecus* monkey D group marker chromosome.

The electrophoretic mobilities of the enzymes G6PD, NP, MDH and
LDH present in an extract prepared from the test cells were comparable to those
of *Cercopithecus* (African green) monkey control cell preparation. No extra
15 bands were observed on the electrophoresis films that might suggest the presence
of cells of a different species in the cell culture. The cells reacted with monkey
antiserum but not with mouse antiserum.

Cells other than those of the submitted culture (Passage 17A, Lot #4706)
were not detected in the culture by either antisera reaction, isoenzyme analysis or
20 by cytogenic evaluation. The chromosome pattern was similar to that previously
observed for cells in the MWCB, Passage 7A, Lot #4214 and specifically related
these cells (Passage 17A, Lot #4706) to the former.

2. Detection of Retroviruses

a. The test article, AGMK Passage 17A, Lot #4706, was analyzed for the presence of types B, C and D retrovirus reverse transcriptase and cellular DNA polymerase activity. The cells were grown in the presence of 30 $\mu\text{g/ml}$

5 bromodeoxyuridine for 24 hours (day 0 to day 1) and then refed with medium without inducer. Culture fluids (unconcentrated and concentrated 20-fold) were harvested on days 2, 3 and 4. No evidence of the presence of retrovirus reverse transcriptase activity was observed in the test article. The results are presented in Table II - A, B and C.

10 b. Detection of Retrovirus Particles by Electron Microscopic

Examination: Cell pellets were prepared from the treated and untreated cultures harvested on day 4 and examined for the presence of virus particles by transmission electron microscopy. No retrovirus-like particles were found in the 200 cells observed.

15 c. Detection of Simian T-Lymphotropic Virus (STLV) in Test Article (AGMK, Passage 17A, Lot #4743) by Polymerase Chain Reaction (PCR)

Technique: The polymerase chain reaction (PCR) provides a method for detection of extremely low numbers of viral nucleic acid sequences, which would not be detectable without amplification. When coupled with a pre-PCR reverse transcriptase step, RNA sequences may be amplified, thus allowing detection of 20 free virus. The protocol used allowed for the amplification of either DNA or RNA derived STLV-specific sequences. The PCR products were tested by

Southern blotting, using chemiluminescent enzyme-linked probe hybridization to the blot.

When tested with the STLV pol primers SK110/SK111, STLV-specific product was not detected in test article reactions, each of which represented 1.0 μ g nucleic acid extracted from the test article. In the positive control series, which contained the equivalent of 0.001 μ g, 0.01 μ g, 0.1 μ g or 1 μ g of nucleic acid extracted from a cell line infected by STLV (STLV_{mn103}), STLV-specific product was detected in all reactions. On the basis of failure to detect reactivity with STLV primers, the test article was judged to be negative for STLV nucleic acid sequences.

d. **Detection of the Simian Immunodeficiency Virus in Test Article (AGMK, Passage 18A, Lot #5506):** Total cellular DNA was extracted from the test article and subsequently tested for the presence of SIV_{agm} virus sequences using primers that amplify a 200 bp fragment of the LTR. This involved PCR with nested sets of primers as follows:

Outer set	#1807 F: CCTCAGAGCTGCATAAAAGCAGAT
	#1808 R: TCACTCAAGTCCCTGTTCGGGCGC
Inner set	#1809 F: TACTAGGAGACCAGCTTGAGCCTG
	#1810 R: TGCTGGAGTTTCTCTCGCCTGGGT

These primers were designed for the conserved U5 and R regions of the LTR and are capable of amplifying all subtypes of SIV_{agm}. The predicted 200 bp fragment was amplified from DNA extracted from a chronically-SIV_{agm}

infected cell line (SIV_{agm}155CEM_{ss}) but the African green monkey cells did not have detectable SIV_{agm} sequences by Southern blot analysis.

3. Tumorigenicity in Athymic Nude Mice

The cells were found to be non-tumorigenic after being clinically
5 evaluated for 150 days. Athymic nude mice were inoculated subcutaneously with approximately 1×10^7 test article cells (AGMK Passage 17A, Lot #4889). Athymic nude mice were similarly inoculated with positive control cells. All 10 positive control inoculated mice had tumors at the site of inoculation. No metastases were observed in the inoculation site (skin), lung, lymph nodes, liver,
10 kidney, spleen or brain of mice inoculated with the test article.

TABLE II-A

Reverse Transcriptase and DNA Polymerase Activities
in Unconcentrated AGMK Passage 17A, Lot #4706

Sample	Reverse Transcriptase ^a (rAdT)		DNA Polymerase ^a (dAdT)	
	Mn ⁺⁺	Mg ⁺⁺	Mn ⁺⁺	Mg ⁺⁺
<i>Test Article Cells^b</i>				
DAY				
2 Untreated	0	0	227	0
Treated	29	0	485	85
3 Untreated	0	0	265	6
Treated	0	0	243	0
4 Untreated	35	0	711	29
Treated	48	0	723	26

^a Expressed as counts per minute ³H-thymidine triphosphate incorporated (mean duplicates minus background).

^b Culture fluids from test article cells.

TABLE II-B

Reverse Transcriptase and DNA Polymerase Activities
in AGMK Passage 17A, Lot #4706
Concentrated 20-fold

Sample	Reverse Transcriptase ^a (rAdT)		DNA Polymerase ^a (dAdT)		
	Mn ⁺⁺	Mg ⁺⁺	Mn ⁺⁺	Mg ⁺⁺	
<i>Test Article Cells^b</i>					
DAY					
2	Untreated	9	0	18	87
	Treated	0	0	15	104
3	Untreated	6	0	32	125
	Treated	9	3	19	136
4	Untreated	22	3	14	114
	Treated	122	2	2,423	516

^a Expressed as counts per minute ³H-thymidine triphosphate incorporated (mean duplicates minus background).

^b Culture fluids from test article cells.

TABLE II-C

Reverse Transcriptase and DNA Polymerase Activities
in AGMK Passage 17A, Lot #4706

Sample	Reverse Transcriptase ^a (rAdT)		DNA Polymerase ^a (dAdT)	
	Mn ⁺⁺	Mg ⁺⁺	Mn ⁺⁺	Mg ⁺⁺
<i>Positive and Negative Controls</i>				
R-MuLV ^c	65,913	2,230	0	16
SMRV ^d	12,031	133,798	1,274	2,825
DNA Polymerase ^e	7,015	57	60,197	6,646
Medium ^f	(163)	(74)	(242)	(107)
Stabilization Buffer ^g	(53)	(59)	(46)	(44)
Medium ^h	(146)	(136)	(321)	(148)

^a Expressed as counts per minute ³H-thymidine triphosphate incorporated (mean duplicates minus background).

^b Culture fluids from test article cells.

^c R-MuLV (Rauscher murine leukemia virus) - Type C virus positive control.

^d SMRV (Squirrel Monkey Retrovirus Preparation) - Type D virus positive control.

^e DNA polymerase - Positive control; prepared from mink lung cells.

^f Medium - Subtracted from unconcentrated, untreated and treated samples.

^g Stabilization buffer - Negative control; subtracted from positive controls and concentrated samples.

^h Medium - Subtracted from R-MuLV and DNA polymerase positive controls.

EXAMPLE 4**Post-Production Testing for Microbial Sterility**

1. **Bacterial Sterility in Fluid Thioglycollate Medium:** Each of 10 culture tubes (9-10 ml medium per tube) was inoculated with 1.0 ml of the AGMK cell suspension (Passage 17A, Lot #4706) containing approx. 5×10^5 cells per ml. Each of 5 tubes was inoculated with 1.0 ml of the original growth medium. Ten cultures were included as uninoculated controls. All cultures were vortex mixed and incubated at $32^\circ\text{C} \pm 2^\circ\text{C}$ for 21 days with observations made on days 1, 3, 6, 8, 10, 13, 15, 17, 20 and 21. No growth was observed in any of the 25 cultures.

2. **Fungal Sterility in Soybean Casein Digest Medium:** Each of 10 culture tubes (9-10 ml medium per tube) was inoculated with 1.0 ml of the AGMK cell suspension (Passage 17A, Lot #4706) containing approx. 5×10^5 cells per ml. Each of 5 tubes was inoculated with 1.0 ml of the original growth medium. Ten cultures were included as uninoculated controls. All cultures were vortex mixed and incubated at $22^\circ\text{C} \pm 2^\circ\text{C}$ for 21 days with observations made on days 1, 3, 6, 8, 10, 13, 15, 17, 20 and 21. No growth was observed in any of the 25 cultures.

3. **Mycobacterial Sterility in Lowenstein-Jensen Egg Medium:** Each of 10 slant culture tubes was inoculated with 0.5 ml of the AGMK cell suspension (Passage 17A, Lot #4706) containing approx. 5×10^5 cells per ml. Each of 10 culture slants was inoculated with 0.5 ml of the original growth medium. Six cultures were included as uninoculated controls. All cultures were incubated at

36°C \pm 1°C horizontally for the first 24 hours and vertically for the remainder of the 8 week observation period with weekly observations for growth. On day 28, one cell suspension inoculated culture was found contaminated with a mold. No growth was observed in any of the remaining 25 cultures after 8 weeks.

5 This assay was repeated with AGMK Passage 18A, Lot #4918, but using 6 culture slants inoculated with the original growth medium. No growth was observed in any of the 22 cultures after 8 weeks. The results of the above Microbial Sterility Assays are summarized in Table III.

4. **Mycoplasma Sterility:** The assay for mycoplasma was performed using
10 both the direct Broth Enrichment and Agar Procedures and the indirect Indicator Cell Culture Procedure with the AGMK cell suspension (Passage 17A, Lot #4706) containing approximately 5×10^5 cells per ml. The cell suspension was found to be negative for mycoplasmas by all procedures.

TABLE III

Microbial Sterility Test Results on the AGMK Cell Suspension -
Passage 17A, Lot #4706 and #4918 at Approximately 5×10^5 Cells/ml

Culture Medium	Vol. Per Culture			Date		Results
	No.	(ml)	Temp.	On Test	Off Test	
<u>Fluid Thioglycollate</u>						
<u>(FIM) LOT VVPL #122-50</u>	10	--	32°C	1/05/93	1/26/93	No Growth
Cell Suspension #4706	10	1.0	(±2°C)			No Growth
Culture Medium	5	1.0		1/05/93	1/26/93	No Growth
<u>Soybean-Casein-Digest</u>						
<u>(SCDM) LOT VVPL #112-51</u>	10	--	22°C	1/05/93	1/26/93	No Growth
Cell Suspension #4706	10	1.0	(±2°C)			No Growth
Culture Medium	5	1.0		1/05/93	1/26/93	No Growth
<u>Lowenstein-Jensen Egg</u>						
<u>Medium - LOT #C3CPPB</u>	6	--	36°C	1/05/93	3/02/93	No Growth
Cell Suspension #4706	10	0.5	(±1°C)			1/10 mold
						on day 28
Culture Medium	10	0.5		1/05/93	3/02/93	No Growth
<u>Lowenstein-Jense Egg</u>						
<u>Medium - LOT #H3CTGX</u>	6	--	36°C	2/23/93	4/20/93	No Growth
Cell Suspension #4918	10	0.5	(±1°C)			No Growth
Culture Medium	6	0.5		2/23/93	4/20/93	No Growth

EXAMPLE 5

**In Vitro Tissue Culture Purity and Safety Tests for
the Detection of Viable Adventitious Microbial Agents**

These assays were performed in roller tube cultures of the following

5 tissue culture systems: Serially passaged African green monkey kidney (AGMK); Primary Human Amnion (PHA); Vero; Primary Rabbit Kidney (PRK); Whole Human Embryo Fibroblast (Fl 5000); and Human Carcinoma of the Larynx (H.Ep-2). Cultures were maintained on Medium EMEM containing 2-10% fetal bovine serum (gamma irradiated) plus antibiotics: gentamicin, 100

10 $\mu\text{g/ml}$; neomycin, 50 $\mu\text{g/ml}$; and fungizone, 2.5 $\mu\text{g/ml}$. Cultures were refed with 2 ml of maintenance medium prior to inoculation.

Testing Procedures: The cell suspensions (Lots #4742/3, #4755 or #4889 containing approximately 5×10^5 cells per ml) were inoculated in 0.5 ml amount per each of 28 roller tubes per tissue culture system. Cultures were incubated at

15 $36^\circ\text{C} \pm 1^\circ\text{C}$ for 14 days with periodic microscopic examination for any cytopathic effect (CPE) and/or cellular degeneration. When necessary to maintain the integrity of the cell films, cultures were refed with 2 ml of fresh maintenance medium. Additional groups of 28 roller tubes per tissue culture system were included as uninoculated controls.

20 On the 7th-8th day of incubation, 6-9 tubes from each series (inoculated and controls) were removed and tested for hemadsorption with 3 red blood cell (RBC) suspensions (guinea pig at 0.1%, chicken at 0.075%, or human at 0.1% in PBS, pH 7.2). Films were rinsed once in the phosphate-buffered saline (PBS)

prior to the addition of 1 ml of the RBC suspension employing 2-3 tubes per RBC. Initially, tubes were incubated at 2° - 8°C for a minimum of 30 minutes and examined microscopically for any signs of hemadsorption. Following manual shaking, tubes were incubated at 35° - 37°C for a minimum of 30 minutes and re-examined microscopically for hemadsorption. All cultures were negative for hemadsorption with all 3 suspensions and at both temperatures. On day 14, after final microscopic examination, the cultures were divided, depending on the specific tissue culture system for final testing. AGMK, PHA and FI5000 Tube Cultures were divided as follows:

1. 9 tubes from each series were tested for hemadsorption with the 3 RBC suspensions as described above for day 7-8;
2. 4 tubes from each series were fixed and stained with a solution of 5% glutaraldehyde + 0.025% crystal violet and, after drying, examined microscopically for any signs of CPE;
3. 8 tubes from each series were challenged with Coxsackie A-9 virus (0.2 ml for each of 2 tubes using a 10^{-2} , 10^{-3} , 10^{-4} or 10^{-5} dilution) for the detection of non-CPE producing agents and/or latent agents via the interference phenomenon; and
4. One (1) tube of each series was included as an uninoculated control for the challenge virus.

Vero, PRK and H.Ep-2 Tube Cultures were divided as follows:

1. 12 tubes from each series were tested for hemadsorption with the 3 RBC suspensions as described above for day 7-8;

2. 7 tubes from each series were fixed and stained with a solution of 5% glutaraldehyde + 0.025% crystal violet and, after drying, examined microscopically for any signs of CPE.

No challenge studies were carried out with the Coxsackie A-9 virus as
5 this agent does not produce any discernible CPE in these cell systems. The results of these *in vitro* tissue culture purity and safety assays are summarized in Tables IV - A to F.

TABLE IV-A

Tissue Culture Purity Tests on the AGMK Cell Suspension -
Passage 17A, LOT #4742/3 at Approximately 5×10^5 Cells/ml

African Green Monkey Kidney Cells - (Passage 12, LOT #4793)

			Day 7 Results		Day 14 Results	
			AGMK Controls		AGMK Controls	
0.1% Guinea Pig RBC	at	2° - 8°C	0/2	0/2	0/3	0/3
	at	35° - 37°C	0/2	0/2	0/3	0/3
0.075% Chicken RBC	at	2° - 8°C	0/2	0/2	0/3	0/3
	at	35° - 37°C	0/2	0/2	0/3	0/3
0.1% Human RBC	at	2° - 8°C	0/2	0/2	0/3	0/3
		35° - 37°C	0/2	0/2	0/3	0/3
Giemsa Stain					0/4	0/4
Coxsackie A-9 Virus Challenge* at	10 ⁻³				2/2	2/2
	10 ⁻⁴				2/2	2/2
	10 ⁻⁵				2/2	2/2
	10 ⁻⁶				2/2	2/2
	Unchallenged Controls				0/1	0/1

* Coxsackie A-9 Challenge results based on a 5-day incubation at 36°C. Prior to challenge, tubes refed with 2 ml of fresh medium.

TABLE IV-B

Tissue Culture Purity Tests on the AGMK Cell Suspension -
Passage 17A, LOT #4742/3 at Approximately 5×10^5 Cells/ml

Primary Human Amnion - (PHA) - (LOT #4720)

			Day 7 Results		Day 14 Results	
			AGMK Controls		AGMK Controls	
0.1 % Guinea Pig RBC	at	2° - 8°C	0/2	0/2	0/3	0/3
	at	35° - 37°C	0/2	0/2	0/3	0/3
0.075 % Chicken RBC	at	2° - 8°C	0/2	0/2	0/3	0/3
	at	35° - 37°C	0/2	0/2	0/3	0/3
0.1 % Human RBC	at	2° - 8°C	0/2	0/2	0/3	0/3
		35° - 37°C	0/2	0/2	0/3	0/3
Giemsa Stain					0/4	0/4
Coxsackie A-9 Virus Challenge* at	10 ⁻³				2/2	2/2
	10 ⁻⁴				2/2	2/2
	10 ⁻⁵				2/2	2/2
	10 ⁻⁶				2/2	2/2
Unchallenged Controls					0/1	0/1

* Coxsackie A-9 Challenge results based on a 5-day incubation at 36°C. Prior to challenge, tubes refed with 2 ml of fresh medium.

TABLE IV-C

Tissue Culture Purity Tests on the AGMK Cell Suspension -
Passage 17A, LOT #4755 at Approximately 5×10^5 Cells/ml

Vero Cells - (Passage 143, LOT #4813)

				Day 7 Results		Day 14 Results	
				AGMK Controls		AGMK Controls	
0.1% Guinea Pig RBC	- at	2° - 8°C	0/3	0/3	0/4	0/4	0/4
	at	35° - 37°C	0/3	0/3	0/4	0/4	0/4
0.075% Chicken RBC	at	2° - 8°C	0/3	0/3	0/4	0/4	0/4
	at	35° - 37°C	0/3	0/3	0/4	0/4	0/4
0.1% Human RBC	at	2° - 8°C	0/3	0/3	0/4	0/4	0/4
		35° - 37°C	0/3	0/3	0/4	0/4	0/4
Giemsa Stain					0/7	0/7	0/7

TABLE IV-D

Tissue Culture Purity Tests on the AGMK Cell Suspension -
Passage 17A, LOT #4889 at Approximately 5×10^5 Cells/ml

Rabbit Kidney Cells - (Primary, LOT #4814)

			Day 7 Results		Day 14 Results*	
			AGMK Controls		AGMK Controls	
0.1% Guinea Pig RBC	at	2° - 8°C	0/4	0/4	0/3	0/3
	at	35° - 37°C	0/4	0/4	0/3	0/3
0.075% Chicken RBC	at	2° - 8°C	0/4	0/4	0/3	0/3
	at	35° - 37°C	0/4	0/4	0/3	0/3
0.1% Human RBC	at	2° - 8°C	0/4	0/4	0/3	0/3
		35° - 37°C	0/4	0/4	0/3	0/3
Giemsa Stain					0/7	0/7

* On day 8, tubes refed with 2 ml of fresh medium.

TABLE IV-E

Tissue Culture Purity Tests on the AGMK Cell Suspension -
Passage 17A, LOT #4889 at Approximately 5×10^5 Cells/ml

Whole Human Embryo Fibroblast Cells - (FL 5000) - (Passage 22, LOT #4842)

			Day 7 Results		Day 14 Results		
			AGMK Controls		AGMK Controls		
0.1 % Guinea Pig RBC	at	2° - 8°C	0/2	0/2	0/2	0/2	
	at	35° - 37°C	0/2	0/2	0/2	0/2	
0.075 % Chicken RBC	at	2° - 8°C	0/2	0/2	0/2	0/2	
	at	35° - 37°C	0/2	0/2	0/2	0/2	
0.1 % Human RBC	at	2° - 8°C	0/2	0/2	0/2	0/2	
		35° - 37°C	0/2	0/2	0/2	0/2	
Giemsa Stain				0/4	0/4		
Coxsackie A-9 Virus Challenge* at				10 ⁻³	2/2	2/2	
				10 ⁻⁴	2/2	2/2	
				10 ⁻⁵	2/2	2/2	
				10 ⁻⁶	2/2	2/2	
Unchallenged Controls				0/1	0/1		

* Coxsackie A-9 Challenge results based on a 4day incubation at 36°C. Prior to challenge, tubes refed with 2 ml of fresh medium.

TABLE IV-F

Tissue Culture Purity Tests on the AGMK Cell Suspension -
Passage 17A, LOT #4755 at Approximately 5×10^5 Cells/ml

Human Carcinoma of the Larynx - H.Ep-2 - (Passage 397, LOT #4808)

			Day 7 Results		Day 14 Results*	
			AGMK Controls		AGMK Controls	
0.1% Guinea Pig RBC	at	2° - 8°C	0/3	0/3	0/4	0/4
	at	35° - 37°C	0/3	0/3	0/4	0/4
0.075% Chicken RBC	at	2° - 8°C	0/3	0/3	0/4	0/4
	at	35° - 37°C	0/3	0/3	0/4	0/4
0.1% Human RBC	at	2° - 8°C	0/3	0/3	0/4	0/4
		35° - 37°C	0/3	0/3	0/4	0/4
Giemsa Stain					0/7	0/7

* On day 10~~8~~, tubes refed with 2 ml of fresh medium.

EXAMPLE 6**In Vivo Tests for the Detection of
Viable Adventitious Microbial Agents**

The following tests were performed employing cell suspensions derived
5 from Passage 17A - (Lots #4706, #4742, #4755 and #4788). These assays were
performed with inocula consisting of approximately 1×10^7 cells per ml
suspended in the culture medium in which they were grown.

1. Tests in Animals

a. **Suckling Mouse Assay:** A total of 10 newborn CD-1 mice randomized
10 from different litters (10 per mother, less than 24 hours old) were inoculated
intracerebrally (I.Cer.) with 0.01 ml and intraperitoneally (I.P.) with 0.1 ml of
the cell suspension from Lot #4706. An additional randomized group of 10
sucklings was inoculated in a similar manner with the original culture medium.
A randomized group of 10 sucklings was included as uninoculated controls. All
15 sucklings were observed daily for 14 days for deaths and/or signs of illness or
distress. On day 2, one (1) suckling from each of the inoculated groups was
found missing and presumed cannibalized. There were no other deaths and none
of the sucklings exhibited any signs of illness or distress over this initial 14 day
observation period.

20 On the 14th day, single pools were prepared of the emulsified tissue
(minus skin and viscera) of the following groups: (a) AGMK inoculated pups
(n=9); (b) culture fluid inoculated pups (n=9); and (c) uninoculated controls
(n=10). A blind passage into newborn CD-1 mice was made of each pool via

the I.Cer. and I.P. routes: each pool into 10 newborns from mixed litters. An additional litter of 10 sucklings was included as uninoculated controls for this blind passage. All sucklings were observed daily for 14 days for deaths and/or signs of illness or distress. There were no deaths recorded for the AGMK or culture medium inoculated pups or for the uninoculated control group; however, on day 3, 4 sucklings were missing and presumed cannibalized in (c) the passaged control group. None of the sucklings exhibited any signs of illness or distress over this final 14 day observation period.

Because none of the AGMK cell suspension or culture medium inoculated sucklings exhibited any signs of a transmissible agent or of Coxsackie virus infection or of any viral infection, and because >90% of these inoculated sucklings (19 of 20) remained healthy and survived the entire observation period, this assay in suckling mice was considered satisfactory.

b. **Adult Mouse Assay:** Each of 20 adult CD-1 mice (VAF, 15-20 grams each) was inoculated I.Cer. and I.P. with 0.03 ml and 0.5 ml, respectively, with the cell suspension Lot #4788. An additional 5 mice were inoculated in a similar manner with the original culture medium. Four mice were included as uninoculated controls. Mice were observed daily for deaths and/or signs of illness or distress over a 21 day period. There were no deaths recorded and all mice survived the entire 21 day observation period with no evidence of lymphocytic choriomeningitis virus infection or any other viral infection. This test in adult mice was considered satisfactory.

- c. **Adult Guinea Pig Assay:** Each of 5 Hartley strain guinea pigs (VAF, 350-450 grams each) was inoculated I.Cer. and I.P. with 0.1 ml and 5.0 ml, respectively, with the cell suspension Lot #4755. An additional guinea pig was inoculated in a similar manner with the original culture medium. One (1) guinea pig was included as an uninoculated control. All were observed daily for deaths and/or signs of illness or distress over a 6 week period; none were recorded. Commencing on day 21, daily rectal temperatures were taken with a digital thermometer and recorded (at approximately 0900 hours) for all guinea pigs until time of sacrifice. The average temperatures (°C) for the AGMK inoculated guinea pigs were: 39.26, 39.30, 39.40, 39.46 and 39.53; for the medium inoculated guinea pig: 39.32; and for the control; 39.37. There were no significant temperature rises indicative of either bacterial or viral infection. All guinea pigs appeared healthy and survived the entire 42-day observation period at which time they were necropsied following euthanasia with Halothane and exsanguination. Inspection of the abdominal and thoracic cavities by the consulting veterinarian did not identify gross lesions or pathological changes. This assay in adult guinea pigs was considered satisfactory.
- d. **Adult Rabbit Assay:** Each of 5 New Zealand white rabbits (1500-2500 gm each) was inoculated intradermally (I.D.) with 0.1 ml per each of 10 sites and subcutaneously (S.Q.) with 9.0 ml of the AGMK cell suspension Lot #4788. One (1) rabbit was similarly inoculated with the original culture medium. One (1) rabbit was included as an uninoculated control. Animals were observed daily for 28 days for deaths, illness, distress and/or lesions at the sites of inoculation.

No deaths were recorded and all rabbits remained healthy without exhibiting any signs of illness or distress or lesions at the sites of inoculation. This test in adult rabbits was considered satisfactory. The results of these *in vivo* animal safety tests are summarized in Table V - A and B.

5 2. Tests in Embryonated Eggs

For these studies, specific pathogen-free embryonated eggs were obtained from SPAFAS, Inc., Reinholds, PA. These eggs came from Flock L061-E.

- a. Allantoic Inoculation: Each of ten 10-day old embryonated eggs was inoculated via the allantoic route with 0.5 ml of the AGMK suspension Lot #4706. An additional 5 eggs were inoculated in a similar manner with the original culture medium. Five (5) eggs were included as uninoculated controls. Eggs were incubated at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 72 hours and then candled for deaths (1 of 5 medium inoculated) and chilled overnight at $2^{\circ}\text{-}8^{\circ}\text{C}$. Allantoic fluids were harvested individually, incubated in a 37°C water bath for 60 minutes to elute any adsorbed agent(s), and then clarified by centrifugation at 2500 rpm for 15 minutes at 20°C . Pools were prepared for each of the 3 sets of eggs employing equal volumes from each individual harvest. The pools were assayed for hemagglutination both undiluted and at a 1:10 dilution with incubation at $2^{\circ}\text{-}8^{\circ}\text{C}$ and at room temperature ($15^{\circ}\text{-}30^{\circ}\text{C}$) using the following erythrocytes in PBS at (pH 7.2): guinea pig at 0.6%; chick at 0.4%; and human at 0.6%. All fluids were negative for hemagglutination at both dilutions and at both temperatures with all 3 RBC suspensions.

The sample pools were subpassaged in 0.5 ml amounts into 10-day old embryonated eggs via the allantoic route as follows: the pool from the AGMK cell suspension into 10 eggs; the pool from the medium into 5 eggs; and the pool from the uninoculated eggs into 5 eggs. An additional 5 eggs were included as uninoculated controls. Eggs were incubated at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 72 hours and then candled for deaths (none recorded) and chilled overnight at $2^{\circ} - 8^{\circ}\text{C}$. Allantoic fluids were harvested, handled and tested as described above. All 4 pools were negative for hemagglutination at both dilutions and at both temperatures with all 3 RBC suspensions.

10 There were no deaths recorded for any of the eggs inoculated with the AGMK cell suspension. Since none of the harvest fluids exhibited any hemagglutination when tested against guinea pig, chick or human RBC, this aspect of the embryonated egg study was considered satisfactory.

b. **Yolk Sac Inoculation:** Each of ten 6-day old embryonated eggs was inoculated into the yolk sac with 0.5 ml of the AGMK cell suspension Lot #4742. Five (5) eggs were similarly inoculated with the original culture medium and 5 eggs were included as uninoculated controls. Eggs were incubated at $36^{\circ}\text{C} [\pm 1^{\circ}\text{C}]$ for 9 days with periodic candling for deaths. Of the medium-inoculated eggs, 2 were found dead (1 each on days 2 & 5); and, of the AGMK-inoculated eggs, 2 were inadvertently broken. No other deaths or losses were recorded. Each group of yolk sacs (AGMK group = 8; medium group = 3; controls = 5) was harvested, pooled, washed in PBS, emulsified by grinding and clarified by centrifugation at 2500 rpm for 20 minutes at 20°C .

The 3 clarified suspensions were subpassaged via the same route into fresh 6-day old embryonated eggs as follows: AGMK group into 10 eggs; medium group into 5 eggs; control group into 5 eggs. An additional 5 eggs were included as uninoculated controls. All eggs were incubated at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 9 days with periodic candling for deaths. On day 7, 3 eggs were found dead; 2 from the AGMK group and 1 from the medium group. There were no other deaths recorded.

This aspect of the embryonated egg study was considered satisfactory, because there was a greater than 80% overall survival (16 of 18) of embryonated eggs (primary and subpassage) inoculated with the AGMK cell suspension and viability of the egg was confirmed.

EXAMPLE 7

Use of Characterized AGMK Cell Substrate
for: (1) Isolation and Growth of Human
Rotaviruses; and (2) Production of
Live Vaccines Containing Such Viruses

The AGMK cells are the only characterized cells tested to date that are suitable for vaccine purposes and support the efficient growth of completely homologous human rotaviruses, i.e., rotaviruses that derive each of their 11 gene segments from a human rotavirus. This is particularly important because a number of candidate homologous human rotavirus vaccine strains are now under development, including those shown in Table V.

As seen in Table V, two candidate human vaccine rotaviruses of G serotype 1 or G serotype 3 recovered from newborn infants were observed to grow to moderately high or high titer in AGMK cells, whereas, each of these neonatal rotavirus strains was markedly restricted in its capacity to grow in two types of cells certified for vaccine production, i.e., FRhL₂ or Vero cells. In addition, one of these neonatal rotavirus strains was also markedly restricted in replicative capacity in human fetal fibroblast cell strain MRC-5 which is licensed for vaccine production.

Similar observations were made with two cold-adapted (*ca*) mutants of human rotavirus of G serotype 1 or G serotype 2 that are also promising candidate live virus vaccine strains. Both human rotavirus mutants grew efficiently in AGMK cells but were significantly or markedly restricted in their growth in certified cells FRhL₂ and Vero.

A. Isolation and Serial Passage of Neonatal Human Rotavirus Strains

Initially, the toxicity of each new preparation of purified trypsin is determined for the AGMK cells to be tested. A known positive stool extract is pre-treated with antibiotics (200 µg/ml gentamicin; 100 µg/ml neomycin; and 5 µg/ml fungizone) for 60 minutes at room temperature. Equal volumes of stool extract and trypsin at pre-determined concentration, e.g., 10 µg/ml, are mixed and allowed to incubate at 37°C in a water bath for 30 minutes. In the interim, the cultures to be inoculated are washed three times with appropriate volumes (e.g., 3 ml for roller tubes, 10 ml for T25 flasks, 30 ml for T75 flasks and 100 ml T175 flasks) of serum-free medium to dilute the serum that had been in the

growth medium in order to prevent components of serum from inactivating trypsin activity.

Roller tube cultures are inoculated with a 0.5 ml volume of the stool/trypsin mixture which is allowed to adsorb at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$ for 60 minutes
5 while rotating on a roller drum apparatus. Cultures are washed once and fed with 3 ml volumes of (EMEM) containing 1% of 10X SPG (sucrose, 2.18M; KH_2PO_4 , 0.038M; K_2HPO_4 , 0.072M; and mono-sodium glutamate, 0.054M) and 1 $\mu\text{g}/\text{ml}$ trypsin. Cell cultures in which trypsin is omitted are also included to control for the effect of trypsin. Incubation is at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$.

10 Cultures are examined microscopically on a daily basis and cultures are harvested when cytopathic effect (CPE) that involves 75 to 100% of the cells is evident. Virus and control cultures are treated similarly with 10% of 10X SPG (v/v) and subjected to one freeze-thaw cycle prior to harvest, sterility testing, distribution for storage, and serial passage. All subsequent passages are carried
15 out either in roller tube cultures or in 25 cm^2 flask cultures employing similar procedures but with variation in trypsin concentration in both pre-treatment and in the final overlay.

An efficient system for producing virus plaques in AGMK cells is developed using methods that are routine in the art so that the virus can be
20 plaque purified in order to obtain a homogeneous population of virus for subsequent vaccine development. The harvest fluids are routinely titrated in the simian MA-104 cells to measure the virus yield obtained. The virus is then triply-plaque-purified in AGMK cells and the resulting clone is amplified by

serial passage in progressively larger tissue culture vessels. The conditions for maximum yield based on trypsin concentration employed in both pre-treatment and final overlay is determined for the progeny of the isolated clone using methods well known in the art.

5 **B. Adaptation and Serial Passage Human Neonatal Rotaviruses or Other
Human Rotavirus Strains**

The human rotaviruses exhibit a very narrow tissue culture host range. As a consequence, isolation and serial passage of various strains are usually performed in commercially available laboratory cell cultures such as simian MA-
10 104 or primary monkey kidney, both of which are unsuitable for vaccine production. The former cells have not been validated as a substrate for vaccine production and the latter cells are notorious for their contamination with various simian microbial agents including retroviruses. In contrast to MA104 or primary monkey kidney cells, cell substrates certified for vaccine production, such as
15 simian FRhL-2 or Vero cells, either fail to support growth of completely homologous human rotaviruses or allow only poor growth.

Many well characterized rotaviruses have been isolated, serially passaged and triply plaque-purified in commercially available laboratory cell culture systems not suitable for vaccine production. In those instances where it was
20 necessary to use such viruses in vaccine development, the viruses were subsequently passaged and biologically cloned in a cell substrate certified for vaccine development and production. Ideally, virus to be used in vaccines should be isolated and passaged only in tissue culture cells certified to be

acceptable for vaccine production. The AGMK cells of this invention meet this requirement because they support efficient growth of fully homologous human rotaviruses (Table V).

To develop a vaccine, routine studies are performed in flask cultures to
5 determine the conditions for maximum yield including trypsin concentration required in both pre-treatment and final overlay. Once adaptation of the vaccine virus to the AGMK cell substrate has been achieved, the virus harvest is treated with ether (1 part ether to 4 parts virus suspension for 60 minutes at room
temperature followed by a 30 minute bubbling with N₂ to drive off ether) to
10 eliminate any ether-sensitive virus(es) that may have been present in the original clinical specimen or acquired subsequently during passage in cell culture prior to initiation of vaccine development.

Live Virus Vaccine and/or Suspension Production

To produce vaccine, it is preferable also to prepare several backup pools
15 of virus as follows:

1. **Primary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared and, after sterility is confirmed, and potency is determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 3rd level
20 back-up to final vaccine production. Total volume may range from 30 ml to 100 ml.
2. **Secondary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) is prepared.

After sterility is confirmed, and potency determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 2nd level back-up to final vaccine production. Total volume may range from 100 ml to 300 ml.

3. **Master Seed Virus Pool or Pre-Production Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) is prepared. After sterility is confirmed, potency is determined, and identity is confirmed by both serology and electrophoretic pattern, the fluids are distributed into multi-sized aliquots with storage at or below -70°C . These pools serve as first level back-up to final production.
- 10 These fluids are subjected to the Tissue Culture Purity (Safety) Testing simultaneously with the vaccine lot. Total volume may range from 500 ml to 1 liter.

4. **Live Virus Vaccine Production:** Volume to be produced is based on the number of doses and containers required plus a minimum of 400 ml of crude fluid for safety testing. In addition to the virus pool, the production of a small pool (200 - 400 ml) of the passaged tissue culture control fluid is recommended for subsequent safety testing together with the crude virus harvest. Culture vessel fluids supplemented with 10% of 10X SPG (v/v) are subjected to one (1) freeze-thaw cycle prior to individual harvest and sterility testing. A sample pool is prepared and assayed for potency. Identity is confirmed by both serology and electrophoretic pattern. The individual harvests are stored at or below -70°C . When sterility, potency and identity are confirmed, crude harvests are thawed and pooled. Samples of about 400 ml are removed for safety testing. The
- 15
- 20

remainder of the fluid is clarified by centrifugation at 1200 g for 20 min. at 5°C, re-pooled, and distributed into final containers.

EXAMPLE 8

5 Adaptation of Human x Animal Reassortant Rotaviruses
 To Growth and Serial Passage in the Characterized
 AGMK Cell Substrate and the Use of Such Cells for
 Production of Live Virus Vaccines

10 Live attenuated rotavirus vaccines can be made using reassortant rotaviruses that derive ten of their eleven gene segments from their animal rotavirus parent and only one gene segment from their human rotavirus parent (usually the gene that encodes the major protective antigen VP7). In one instance, human x rhesus monkey rotavirus reassortants are in a late stage of vaccine development and an application for a license has been submitted to the FDA.

15 All vaccine candidate human X animal reassortant rotaviruses studied to date have exhibited a very broad tissue culture host range attributable to the animal rotavirus parent, which contributes ten of its eleven gene segments to the reassortant. Although these vaccine candidate reassortments can proliferate to high titer in other cell lines certified for vaccine production, comparative studies
20 with AGMK cell substrate indicate that the AGMK cell substrate provided a significant advantage in viral growth yield over FRhL2 cells in every instance tested (Table V). In addition, the AGMK cell substrate has proved to be

superior to Vero cells for the growth of human x rhesus rotavirus reassortants as well as rhesus rotavirus itself (Table V). In these instances, RRV and its human rotavirus reassortants grew in the AGMK cell line to a titer that was 16- to 100-fold higher than in FRhL2 cells and 10- to 1200-fold higher than in Vero cells (Table V). The only instances in which the AGMK cell substrate was not superior to Vero cells involved the human x bovine rotavirus reassortants where viral growth yield in the two cell lines was equivalent (Table V).

In addition to yield, the AGMK cell substrate has other advantages: (a) pre-treatment of virus with trypsin is not required; (b) the concentration of trypsin required in culture medium is reduced; (c) there is a significant reduction in the amount of virus required in the inoculum; (d) incubation time is shortened; and (e) the infected cell monolayer is completely lysed. Vaccine production is carried out as outlined above.

EXAMPLE 9

Adaptation of Human Astrovirus to Growth, Serial Passage and Subsequent Live Virus Vaccine Production in the Characterized AGMK Cell Substrate

Previously, all human astroviruses have been isolated or propagated in cells inappropriate or unacceptable for vaccine production. A Type 2 strain of human astrovirus was successfully adapted to growth and serial passage in the AGMK cell substrate from seed virus grown in the simian LLCMK₂ cell line. Flask cultures of AGMK cell grown virus were studied to determine the optimal

conditions for maximum yield based on trypsin concentrations employed in both pre-treatment and final overlay. Once adaptation was achieved, the virus harvest was treated with ether (1 part ether to 4 parts virus suspension for 60 minutes at room temperature followed by a 30 minute bubbling with N₂ to drive off ether) to eliminate any ether sensitive simian virus that may have been picked up during passage and plaquing in the commercially available laboratory cell culture systems. Fluid production was similar to that outlined for the human rotavirus.

EXAMPLE 10

Propagation of Live Attenuated Poliovirus Vaccine Strains in the Characterized AGMK Cell Substrate

The Sabin live attenuated oral poliovirus vaccine strains were grown in the AGMK cells at passages 15 and 16. Each of the vaccine strains grew to high titer in AGMK cells (type 1 virus $10^{8.3}$ TCID₅₀/ml, type 2 virus $10^{8.1}$ TCID₅₀/ml, and type 3 virus $10^{8.6}$ TCID₅₀/ml). These titers translate into 100 to 1000 vaccine doses per ml of tissue culture harvest and are equivalent to the amount of virus produced by licensed primary AGMK cells and MRC-5 cells.

EXAMPLE 11

Isolation and/or Adaptation of Enteroviruses to Growth, Serial Passage and Subsequent Live Virus Suspension and/or Vaccine Production in the Characterized AGMK Cell Substrate

Enteroviruses such as Coxsackie A9 virus were adapted to growth and serial passage in the characterized AGMK cells. These cells are used to produce live virus suspensions and vaccine. Several seed virus pools have been produced.

5 Live Virus Vaccine and/or Suspension Production

1. **Primary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared and after sterility is confirmed and potency is determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 3rd level back-up to
10 final vaccine production. Total volume may range from 30 ml to 100 ml.
2. **Secondary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared. After sterility is confirmed, and potency determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 2nd level back-up to
15 final vaccine production. Total volume may range from 100 ml to 300 ml.
3. **Master Seed Virus Pool or Pre-Production Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared. After sterility is confirmed, potency is determined, and identity confirmed by serology. The fluids are distributed into
20 multi-sized aliquots with storage at or below -70°C. These pools serve as first level back-up to final production. These fluids are subjected to the Tissue Culture Purity (Safety) Testing simultaneously with the vaccine lot. Total volume may range from 500 ml to 1 liter.

4. **Live Virus Vaccine Production:** Volume to be produced is based on the number of doses and containers needed plus a minimum of 400 ml of crude fluid for the required safety testing. In addition to the virus pool, the production of a small pool (200 - 400 ml) of the passaged tissue culture control fluid is
- 5 recommended for subsequent safety testing together with the crude virus harvest. Culture vessel fluids supplemented with 10% of 10X SPG (v/v) are subjected to one (1) freeze-thaw cycle prior to individual harvest and sterility testing. A sample pool is prepared and assayed for potency and identity is confirmed by serology. The individual harvests are stored at or below -70°C. When sterility,
- 10 = potency and identity are confirmed, crude harvests are thawed and pooled. Samples of about 400 ml are removed for safety testing. The remainder of the fluid is clarified by centrifugation at 1200 g for 20 min. at 5°C, re-pooled, and distributed into final containers.

EXAMPLE 12

- 15 **Isolation and Adaptation of Respiratory Syncytial Virus (RSV) to Growth and Serial Propagation in the Characterized AGMK Cell Substrate and Use of These Cells for Production of a Live Virus Vaccine**

- Respiratory syncytial viruses, subgroup A and B types, have been readily
- 20 isolated in these AGMK cell cultures directly from human nasal and/or throat secretions. Specimens are centrifuged at 3000 rpm for 10 minutes to remove any articulate matter and then treated with antibiotics (200 µg/ml gentamicin; 100 µg/ml neomycin; and 5 µg/ml fungizone) for 60 minutes at room temperature.

The antibiotic-treated specimens are inoculated into roller tube cultures of the AGMK cells and incubated at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$ on a roller drum. Once cytopathic effects are evident, cultures are harvested and further passages are carried out either in roller tubes or in flask cultures with subsequent serial passage and plaque purification. Subsequently, virus mutants are produced by chemical mutagenesis or are selected by passage at suboptimal temperature using the AGMK cells in either instance. Live virus vaccine is then prepared using the AGMK cells.

Final viral yield of candidate live attenuated RSV vaccine strains grown in the AGMK cells was greater than that of these viruses grown in Vero cells, which were previously the most efficient cell substrate certified for production of RSV vaccines. An example is seen in Table VI, where the virus yield of a cold-passaged, temperature sensitive mutant of RSV (designated RSV A2 *cpts248/404*) is shown to be 4- to 16-fold higher in the AGMK cell line than in Vero cells. Also, the amount of RSV produced by the human fetal diploid cell line MRC-5, which has been licensed for use in production of virus vaccines, is 10- to 100-fold less than in AGMK cells. This mutant is highly attenuated but still capable of inducing resistance to RSV in susceptible chimpanzees, which represent the most relevant animal surrogate for evaluation of attenuation and protective efficacy of RSV mutants prior to initiation of clinical trials in humans.

Live attenuated RSV vaccines are designed to be used in very young infants, the population at greatest risk of life-threatening RSV disease. Because of the need to immunize such young individuals, the live vaccine must be very

attenuated (i.e., demonstrably weakened). Thus, it is noteworthy that the RSV A2 *cpts248/404* mutant was able to attain a high titer in AGMK cells. Clearly, success of a vaccine containing this mutant and live RSV vaccines in general will require attainment of a high level of viral yield.

- 5 To produce live virus vaccine, conditions for good growth and viral yield, such as temperature of incubation and inoculum size, are determined by routine methods with production proceeding as follows:

Live Virus Vaccine and/or Suspension Production

1. **Primary Seed Virus Pool:** A pool of virus and serially passaged tissue
10 culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared and after sterility is confirmed and potency is determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 3rd level back-up to final vaccine production. Total volume may range from 30 ml to 100 ml.
2. **Secondary Seed Virus Pool:** A pool of virus and serially passaged tissue
15 culture control fluids supplemented with 10% of 10X SPG (v/v) is prepared. After sterility is confirmed, and potency determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 2nd level back-up to final vaccine production. Total volume may range from 100 ml to 300 ml.
3. **Master Seed Virus Pool or Pre-Production Seed Virus Pool:** A pool of
20 virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared. Sterility is confirmed, potency is determined, and identity confirmed by serology. The fluids are distributed into multi-sized aliquots with storage at or below -70°C. These pools serve as first level back-up

to final production. These fluids are subjected to the Tissue Culture Purity (Safety) Testing simultaneously with the vaccine lot. Total volume may range from 500 ml to 1 liter.

4. **Live Virus Vaccine Production:** Volume to be produced is based on the number of doses and containers needed plus a minimum of 400 ml of crude fluid for the required safety testing. In addition to the virus pool, the production of a small pool (200 - 400 ml) of the passaged tissue culture control fluid is recommended for subsequent safety testing together with the crude virus harvest. Culture vessel fluids supplemented with 10% of 10X SPG (v/v) are harvested directly without undergoing a freeze - thaw cycle. A sample pool is prepared and assayed for potency and identity is confirmed by serology. The crude harvests are pooled and aliquots of about 400 ml are removed for safety testing. The remainder of the fluid is clarified by centrifugation at 1200 g for 20 min. at 5°C, re-pooled, and distributed into final containers.

15

EXAMPLE 13

Isolation and Adaptation of Parainfluenza Viruses to Growth and Serial Propagation in the Characterized AGMK Cell Substrate and Use of These Cells for Production of a Live Virus Vaccine

- Parainfluenza viruses types 1, 2 and 3 were isolated previously in primary AGMK cell cultures directly from human nasal and/or throat specimens. Subsequently, virus types 1, 2 and 3 were readily adapted to growth in the characterized AGMK cells. For isolation in the characterized AGMK cell

substrate clinical specimens are centrifuged at 3000 rpm for 10 minutes to remove any particulate matter and then treated with antibiotics (200 µg/ml gentamicin; 100 µg/ml neomycin; and 5 µg/ml fungizone) for 60 minutes at room temperature. The antibiotic-treated specimens are inoculated into the

5 AGMK cell substrate roller tube cultures and incubated at $36^{\circ}\text{C} \pm 1^{\circ}\text{C}$ on a roller drum. Once cytopathic effects and/or hemadsorption (with 0.1% guinea pig RBC in PBS, pH 7.2) is evident, cultures are harvested with further passages carried out either in roller tubes or in flask cultures with subsequent serial passage followed by plaque purification. Subsequently, virus mutants are

10 produced by chemical mutagenesis or are selected by passage at suboptimal temperature, using the AGMK cell substrate in either instance.

An attenuated cold-adapted (*ca*) mutant of parainfluenza virus type 3 (designated PIV3 *ca* cold passage 45) was successfully grown to high titer in the AGMK cell substrate. Growth yield of the mutant was similar to the yield

15 obtained in Vero cells, a certified simian cell line that has not yet been licensed for production of live virus vaccine for use in humans. However, as shown in Table VII, final viral yields in the AGMK cells were approximately 30-fold higher than in the commonly used, certified fetal rhesus lung cell strain, FRhL-2, which also has not been licensed as yet for production of human viral vaccines.

20 The PIV3 *ca* cold passage 45 mutant, which is now in Phase II clinical trials, is satisfactorily attenuated for young infants who have not been previously infected with naturally occurring PIV3. Although the mutant is very attenuated in young infants, it nonetheless stimulates production of protective levels of

serum PIV3 antibodies. Based upon the results of clinical trials of the *ca* mutant in young infants, we calculate that 1 ml of the viral yield from infected AGMK cells contains 30 vaccine doses of PIV3 *ca* cold passage 45. Attainment of this level of viral growth during vaccine production is particularly noteworthy

- 5 because the live PIV3 vaccine has been weakened so that it can be administered safely to infants and young children, the target population for immunization.

To produce live virus vaccine, conditions for good growth and viral yield, such as temperature of incubation and inoculum size, are determined by routine methods with production proceedings as follows:

10 **Live Virus Vaccine and/or Suspension Production**

1. **Primary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared and, after sterility is confirmed and potency is determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 3rd level
15 back-up to final vaccine production. Total volume may range from 30 ml to 100 ml.
2. **Secondary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) is prepared. After sterility is confirmed, and potency determined, the fluids are distributed
20 into small aliquots for storage at or below -70°C to serve as 2nd level back-up to final vaccine production. Total volume may range from 100 ml to 300 ml.
3. **Master Seed Virus Pool r Pre-Production Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10%

of 10X SPG (v/v) are prepared. Sterility is confirmed, potency is determined, and identity confirmed by serology. The fluids are distributed into multi-sized aliquots with storage at or below -70°C. These pools serve as first level back-up to final production. These fluids are subjected to the Tissue Culture Purity

5 (Safety) Testing simultaneously with the vaccine lot. Total volume may range from 500 ml to 1 liter.

4. **Live Virus Vaccine Production:** Volume to be produced is based on the number of doses and containers needed plus a minimum of 400 ml of crude fluid for the required safety testing. In addition to the virus pool, the production of a
10 small pool (200 - 400 ml) of the passaged tissue culture control fluid is recommended for subsequent safety testing together with the crude virus harvest. Culture vessel fluids supplemented with 10% of 10X SPG (v/v) are subjected to one (1) freeze-thaw cycle prior to individual harvest and sterility testing. A sample pool is prepared and assayed for potency and identity. The individual
15 harvests are stored at or below -70°C. When sterility, potency and identity are confirmed, crude harvests are thawed and pooled. Samples of about 400 ml are removed for safety testing. The remainder of the fluid is clarified by centrifugation at 1200 g for 20 min. at 5°C, re-pooled, and distributed into final containers.

20

EXAMPLE 14

**Isolation and/or Adaptation of Influenza Viruses to Growth
in the Characterized AGMK Cell Substrate and Use of These Cells
for Production of a Live or Inactivated Virus Vaccine**

Influenza viruses type A, subtypes H3N2, and H1N1 and type B, have been readily adapted to growth in this characterized AGMK cell line. A significant advantage of isolating and growing influenza viruses in a mammalian cell line rather than in embryonated chicken eggs stems from the fact that the hemagglutinin (the major protective viral protein of influenza virus) retains the same amino acid sequence and conformation as that present in a virus isolated from humans. In contrast, mutants bearing mutations affecting the hemagglutinin are selected during adaptation of virus to growth in eggs. These mutations confer upon the mutant a growth advantage in eggs.

10 The advantages of producing influenza virus vaccines, either live or killed, in a mammalian tissue culture system rather than in embryonated eggs derive from retention of the major protective antigenic sites on the influenza virus hemagglutinin (HA) when virus is grown in mammalian tissue culture. Adaptation of influenza A or B virus to efficient growth in embryonated chicken
15 eggs frequently results in selection of viral mutants that possess mutations affecting one of the major protective antigenic sites on the viral HA. Loss of such an HA epitope that induces protective antibodies diminishes the effectiveness of a vaccine containing this form of viral antigenic mutant. A number of studies have shown that influenza viruses isolated and propagated only
20 in mammalian cell culture are more closely related antigenically to naturally circulating viruses than are egg-adapted influenza virus strains. For this reason it would be advantageous to isolate and maintain influenza A and B virus vaccine strains in a mammalian cell culture system, preferably primate, so as to maintain

authentic antigenic structure of HA which is the major protective protein of these viruses.

As shown in Table VIII, a variety of influenza A viruses of subtype H1N1 or H3N2 grew efficiently in the AGMK cell substrate. Various influenza B virus strains also grow in the AGMK cells, albeit somewhat less well than the influenza A viruses. Since egg-propagated viruses were used as the inoculum in this experiment, the viral yields shown in Table VIII are minimal estimates because egg adaptation would be expected to select for mutants able to grow efficiently in eggs. Such mutants often grow less well in primate cells than the original naturally occurring virus as a consequence of the mutations responsible for egg adaptation.

The AGMK cells can be used to isolate influenza A and B viruses from infected humans and serve as a cell substrate for production of virus that is used to produce an inactivated vaccine. Alternatively, the AGMK cells can be used to isolate and propagate virus during the derivation of attenuated mutants that are used for formulation of a live attenuated virus vaccine.

EXAMPLE 15

Isolation and/or Adaptation of Human and Simian Hepatitis A Viruses to Growth, Serial Passage and Subsequent Live and Inactivated Virus Vaccine Production in the Characterized AGMK Cell Substrate

Wild-type hepatitis A virus strains of human origin grow poorly or not at all in cell culture on primary isolation: they must be adapted to grow efficiently

in cell culture by the laborious procedure of serial passage. In contrast, wild-type HAV of simian origin replicates moderately well during primary isolation in certain primate cells. However, certified or licensed cell culture systems suitable for primary isolation and adaptation of HAV strains of human or simian origin
5 are not readily available or do not support the replication of HAV to high levels. Although the examples are for human HAV, it will be obvious to one skilled in the art that the techniques can be applied to HAV strains of simian origin.

Human hepatitis A virus, strain HM-175 (HAV/7), a strain suitable for a candidate live attenuated vaccine as well as a candidate inactivated vaccine, was
10 compared for ability to replicate in AGMK cells, FRhK-4 (clone 11-1, the most sensitive cell line for replicating this virus to date), and licensed MRC-5 human diploid cells. Conditions for cell culture were those commonly used for the replication of HAV in FRhK-4 cells.

As seen in Table IX and Figure 3, HAV replicated to the same titer and
15 at the same rate in the AGMK cell substrate, which is suitable for vaccine development and production, as in FRhK-4, clone 11-1 cells, which are not suitable for vaccine development. Furthermore, the size of the HAV plaques, as measured by the radioimmunofocus assay, was the same in both cell types. In contrast, HAV strain HM-175 (HAV/7) did not replicate efficiently in MRC5
20 cells (a cell strain licensable for vaccine development) without extensive adaptation by serial passage.

Live Virus Vaccine and/ r Suspension Production

1. **Primary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared and after sterility is confirmed and potency is determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 3rd level back-up to
5 final vaccine production. Total volume may range from 30 ml to 100 ml.
2. **Secondary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) are prepared. After sterility is confirmed and potency determined, the fluids are distributed into small aliquots for storage at or below -70°C to serve as 2nd level back-up to
10 final vaccine production. Total volume may range from 100 ml to 300 ml.
3. **Master Seed Virus Pool or Pre-Production Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG (v/v) is prepared. Sterility is confirmed, potency is determined, and identity confirmed by serology. The fluids are distributed into multi-sized
15 aliquots with storage at or below -70°C. These pools serve as first level back-up to final production. These fluids are subjected to the Tissue Culture Purity (Safety) Testing simultaneously with the vaccine lot. Total volume may range from 500 ml to 1 liter.
4. **Live Virus Vaccine Production:** Volume to be produced is based on the
20 number of doses and containers needed plus a minimum of 400 ml of crude fluid for the required safety testing. In addition to the virus pool, the production of a small pool (200 - 400 ml) of the passaged tissue culture control fluid is recommended for subsequent safety testing together with the crude virus harvest.

Culture vessel fluids supplemented with 10% of 10X SPG (v/v) are subjected to one (1) freeze-thaw cycle prior to individual harvest and sterility testing. A sample pool is prepared and assayed for potency and identity. The individual harvests are stored at or below -70°C. When sterility, potency and identity are confirmed, crude harvests are thawed and pooled. Samples of about 400 ml are removed for safety testing. The remainder of the fluid is clarified by centrifugation at 1200 g for 20 min. at 5°C, re-pooled, and distributed into final containers.

Inactivated Virus Vaccine and/or Suspension Production

- 10 1. **Primary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG [v/v] that are sterility confirmed, potency determined and distributed into small aliquots for storage at -70°C, or below, to serve as 3rd level back-up to final vaccine production. Total volume may range from 30 ml to 100 ml.
- 15 2. **Secondary Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG [v/v] that are sterility confirmed, potency determined and distributed into small aliquots with storage at -70°C, or below, to serve as 2nd level back-up to final vaccine production. Total volume may range from 100 ml to 300 ml.
- 20 3. **Master Seed Virus Pool or Pre-Production Seed Virus Pool:** A pool of virus and serially passaged tissue culture control fluids supplemented with 10% of 10X SPG [v/v] that are sterility confirmed, potency determined, identity confirmed, and distributed into multi-sized aliquots with storage at -70°C, or

below. Pools serve as 1st level back-up to final production. Fluids to be subjected to the Tissue Culture Purity (Safety) Testing simultaneously with the vaccine lot. Total volume may range from 500 ml to 1 liter.

4. **Inactivated Virus Vaccine Production:** Volume to be produced is based
- 5 on the number of doses and containers contracted for plus a minimum of 400 ml of crude fluid needed for the required safety testing. In addition to the virus pool, the production of a small pool [200 - 400 ml] of the passaged tissue culture control fluid is recommended for subsequent safety testing together with the crude virus harvest. Culture vessel fluids supplemented with 10% of 10X SPG
- 10 [v/v] are subjected to one (1) freeze-thaw cycle prior to individual harvest and sterility testing. A sample pool is prepared and assayed for potency and identity and the individual harvests stored at -70°C, or below. When sterility, potency and identity are confirmed, crude harvests are thawed, pooled, clarified and purified before inactivation with formalin under standard conditions. The
- 15 vaccine is then distributed into final containers and safety-tested for sterility.

TABLE V

Comparison of Growth Patterns of Candidate
Rotavirus Vaccines in Various Cell Cultures

Candidate Rotavirus Vaccine Strain (Serotype) ^b	Titer of Indicated Candidate Live Vaccine Virus Strain Produced in Indicated Cell Culture as Determined by Plaque Assay in MA104 Cells (log ₁₀ pfu/ml) ^a			
	AGMK ^c	FRhL-2	Vero ^c	MRC5 ^c
<i>Human Strains</i>				
D - ca30°C (G1;P1A) ^d	6.1	No Growth	4.7	NT ^e
DS-1 - ca30°C (G2;P1B) ^d	6.3	No Growth	No Growth	NT
M37 - neonatal (G1;P2)	5.2	No Growth	No Growth	NT
Neonatal (G3;P2)	5.8	Inconsistent, poor growth	Inconsistent, poor growth	No Growth
<i>Human x Bovine Viral Reassortants^f</i>				
D x UK (G1;P7)	7.1	5.8	7.1	NT
DS-1 x UK (G2;P7)	7.3	5.3	7.0	NT
P x UK-2 (G3;P7)	7.0	5.4	6.9	NT
ST-3 x UK-2 (G4;P7)	7.2	5.8	6.6	NT
<i>Rhesus Rotavirus or Human x Rhesus Monkey Viral Reassortants^f</i>				
D x RRV (G1;P5B)	7.9 ^g	6.7	6.6 ^g	NT
	7.9		5.7	
	8.3		5.1	
DS-1 x RRV (G2;P5B)	7.6 ^g	5.7	6.6 ^g	NT
	7.6		6.1	
	7.7		4.8	
RRV-2 (G3;P5B)	8.5 ^g	6.7	7.2 ^g	NT
	8.6		7.3	
	8.7		6.4	
ST3 x RRV (G4;P5B)	7.6 ^g	6.1	6.3 ^g	NT
	7.8		5.8	
	7.7		4.6	

^a Virus suspension grown in indicated cells.

^b Serotype denotes viral outer capsid proteins (G and P) that induce protective neutralizing antibodies.

^c AGMK = African green monkey kidney cells (derived from *Cercopithecus ethiopus* monkey); FRhL-2 = Fetal rhesus-lung cells; MRC5 = fibroblastic cells derived from fetal human lung. FRhL-2 and MRC5 are semi-continuous cell strains and can not be used beyond passage 32 for vaccine production; Vero is a continuous cell line; the number of times AGMK cells can be passaged serially and remain suitable for use in vaccine development has not been determined yet but is at least 16.

^d Cold-adapted (ca) mutants of human rotavirus selected by growth and serial passage in cell culture at suboptimal temperature that does not allow efficient growth of naturally occurring (i.e., wild type) human rotavirus.

^e NT = not tested.

^f A single gene (encoding G serotype) derived from human rotavirus and all the remaining 10 genes derived from bovine or rhesus rotavirus.

^g Growth yield from second passage in AGMK or Vero cells of virus previously passaged in FRhL-2 cells; values shown indicate growth yield from cell culture inoculated with 1:10; 1:100 or 1:1000 dilution of first passage material, respectively.

Note: AGMK cells were used at passages 10, 11, 14 or 15.

TABLE VI

Efficient Replication of a Candidate Live Attenuated RSV Vaccine Strain
(RSV A2 *cpts248/404*) in AGMK Substrate

Vero Cell Grown RSV A2 <i>cpts248/404</i> Used to Initiate Infection in	Quantity of Virus ($\text{Log}_{10}\text{pfu/ml}$) Produced in Cell Cultures Inoculated With*	
	1 ml	5 ml
Vero Cells	5.80	5.59
AGMK Cells	6.43	6.80

* RSV A2 *cpts248/404* mutant grown and titrated at 32°C. Titrations performed in HEp2 cells at 32°C.

Note: AGMK -cells used at passages 14 and 15.

TABLE VII

Efficient Replication of a Candidate Live Attenuated Parainfluenza Virus
Type 3 Vaccine Strain (PIV3 Cold-Adapted Cold Passage 45) in
AGMK Cell Substrate

Temperature of Incubation of PIV3 <i>ca</i> cp45	Cell Substrate Used for Virus Growth	Quantity of Virus Produced ^b (Log ₁₀ pfu/ml)	Hemagglutinin Titer (Reciprocal)
32°C	AGMK	7.37	256
	Vero	7.05	256
30°C	AGMK	7.49	256
	Vero	7.22	128
26°C	AGMK	7.54	256
	Vero	7.78	512

^a Inoculum was FRhL2 cell grown virus currently being evaluated in clinical trials. The titer of this FRhL2 cell virus suspension was 10^{6.0}pfu/ml.

^b Inoculated cell cultures harvested after 14 days of incubation except 32°C AGMK cultures which were harvested at 7 days because of accelerated development of viral cell destructive effects.

Note: AGMK cells used at passages 13 and 14.

TABLE VIII

Growth of Human Influenza A and B Viruses in AGMK Cell Substrate

Type and Subtype	Virus Strain	Number of Passages in AGMK Cells at 36°C ^a	Quantity of Virus Produced at the highest passage level as Determined by Titration in Indicated Cell Substrate	
			AGMK	MDCK ^b
A H3N2	A/Wash/897/80	1	6.05	6.60
	A/Korea/1/82	3	7.15	7.79
	A/Bethesda/1/85	2	6.48	6.90
	A/Los Angeles/2/87	3	7.16	7.76
	A/Shandong/9/93	3	7.28	7.78
A H1N1	A/Texas/1/85	1	5.31	4.87
	A/Kawasaki/9/86	3	6.50	6.81
	A/Texas/36/91	1	5.23	6.04
B	B/Texas/1/84	1	<3.00	4.78
	B/Ann Arbor/1/86	1	3.78	5.58
	B/Victoria/2/87	1	<3.00	4.82
	B/Yamagata/16/86	1	4.41	5.72

^a Passage in AGMK cell substrate initiated with egg grown influenza virus at a dilution of 10^{-3} or 10^{-4} . The inoculum for passage in cell culture was also 10^{-3} or 10^{-4} (except B/Victoria/2/87 which was 10^{-2}) in order to minimize generation of defective, interfering virus particles. Tissue culture medium and agarose overlay in plaque assays contained 0.5 μ g/ml of trypsin.

^b MDCK is a continuous cell line of canine kidney cells often used for titration of human influenza viruses.

TABLE IX

Ability of AGMK Cells to Support Efficient Growth and Plaque
Formation of Candidate Hepatitis A Virus Vaccine Strain HM-175 (HAV/7)*

HAV Property	Cell Culture Used for Virus Replication		
	AGMK ^a	FRhK-4(11-1) ^a	MRC-5 ^a
Titer (log ₁₀ pfu/ml) of Virus Achieved in Inoculated Culture ^b	7.4	7.4	Little or No Growth ^c
Plaque Size	Large	Large	---

* Candidate strain for both inactivated and live attenuated HAV vaccines.

^a AGMK = African green monkey kidney derived from Freeze #2129. FRhK-4(11-1) = HAV-permissive cell clone derived from fetal rhesus kidney-4 cells (Funkhouser AW, Purcell RH, D'Hondt E, Emerson SU, Attenuated hepatitis A virus: Genetic determinants of adaptation to growth in MRC-5 cells. *J Virol* 68:148-157; 1994). MRC-5 = fetal human lung fibroblast cell substrate.

^b Determined by plaque assay on AGMK and FRhK-4(11-1) cells.

^c Growth not detected by plaque assay performed with FRhK-4(11-1) cell culture.

Note: AGMK cells used at passages 12, 13, 15 and 16.

WHAT IS CLAIMED IS THE FOLLOWING:

1. Serially passaged cells derived from the kidneys of an African
Green Monkey that are substantially free of viable adventitious microbial agents.
2. A cell substrate which, according to claim 1, wherein said cell
5 substrate is an aneuploid female *Cercopithecus* monkey cell substrate with
chromosome counts in the diploid range.
3. A cell line according to claim 1 or 2, which is capable of
supporting the growth of a large number of different viruses.
4. Cell substrate ATCC # CRL 11756, and mutants and variants
10 thereof.
5. A method for isolating a virus in the cell substrate of claim 1
comprising the steps of:
 - i) inoculating said cells with a sample containing virus;
 - ii) incubating said inoculated cells to permit proliferation of
15 said suspension of virus; and
 - iii) harvesting said virus.

6. A method, according to claim 5, wherein the virus is a human rotavirus.
7. A method according to claim 5, wherein said virus is a human X animal rotavirus reassortant.
- 5 8. A method, according to claim 5, wherein the virus is a human astrovirus.
9. A method, according to claim 5, wherein the virus is an enterovirus.
- 10 10. A method according to claim 5, wherein the virus is a Sabin live attenuated oral polio virus vaccine strain.
11. A method, according to claim 5, wherein the virus is a respiratory virus.
12. A method, according to claim 5, wherein the virus is a hepatitis A virus.
- 15 13. A rotavirus produced according to the method of claim 6.

14. An astrovirus produced according to the method of claim 8.
15. An enterovirus produced according to the method of claim 9.
16. A Sabin live attenuated oral polio virus produced according to the method of claim 10.
- 5 17. A respiratory virus produced according to claim 11.
18. A hepatitis A virus produced according to claim 12.
19. A method of producing a vaccine comprising the steps of:
- i) producing a virus by the method of claim 5; and
 - ii) using said virus to prepare a vaccine.
- 10 20. A method, according to claim 19, wherein the virus is a rotavirus.
21. A method, according to claim 19, wherein the virus is a human rotavirus.
22. A method, according to claim 19, wherein the virus is a human X animal rotavirus reassortant.

23. A method, according to claim 19, wherein the virus is a human
astrovirus.

24. A method, according to claim 19, wherein the virus is an
enterovirus.

5

25. A method according to claim 19, wherein the virus is a Sabin live
attenuated oral poliovirus.

26. A method, according to claim 19, wherein the virus is a
respiratory virus.

10 27. A method, according to claim 19, wherein the virus is a hepatitis
A virus.

1/3

Disassociated Kidneys Freeze #2129

Primary Flasks - Freeze #2076 = MSCB of 1/29/91

p1 - L#3704

p2 - L#3738

p3 - L#3784

p4 - L#3901

p5A - L#4019 frozen 8/3/92 as L#4098 = MWCB

FIG. 1

2/3




FIG. 2

Radioimmunofocus Assay of HAV/7: Time Course for Plaque Formation




FIG. 3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15118

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 39/12, 39/13, 39/15, 39/145, 39/29; C12N 5/06, 7/02

US CL : 424/204.1, 205.1, 209.1, 215.1, 217.1, 226.1; 435/235.1, 240.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/204.1, 205.1, 209.1, 215.1, 217.1, 226.1; 435/235.1, 240.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)


Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X ---- A	US, A, 4,040,905 (PETRICCIANI ET AL.) 09 August 1977, see entire document.	1, 3, 5, 9-11, 15-17, 19, 24- 26 ----- 2, 4
X	US, A, 4,571,385 (GREENBERG ET AL.) 18 February 1986, see entire document, especially column 4.	1, 3, 5-7, 13, 19-22
X	Journal of General Virology, Volume 57, issued 1981, LEE ET AL., "Serial Propagation of Astrovirus in Tissue Culture with the Aid of Trypsin", pages 421-424, see entire document.	1, 3, 5, 8, 14, 19, 23
X	US, A, 4,783,407 (PROVOST ET AL.) 08 November 1988, see entire document.	1, 3, 5, 12, 18- 19, 27

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"E" earlier document published on or after the international filing date	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"A" document member of the same patent family
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 29 FEBRUARY 1996	Date of mailing of the international search report 22 MAR 1996
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer TERRY A. MCKELVEY  Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15118

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Abstracts of the Annual Meeting of the American Society for Microbiology, 23rd Annual Meeting, issued March 1983, PRICE ET AL, "The Use of Serially Propagated Monkey Kidney Cell Cultures for the Preparation of High-Titered Hemagglutinin and/or Complement-Fixing Antigens of the Human Parainfluenza Viruses-Types 1, 2 and 3", page 299, column 2, abstract T22.	1, 3, 5, 11, 17, 19, 26
X, P	Biologicals, Volume 23, Number 3, issued September 1995, HORAUD, "Viral Vaccines and Residual Cellular DNA", pages 225-228, see page 227, column 1.	1, 3

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15118

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, DIALOG

search terms: african green monkey, kidney, agmk, vero, bac, cercopithecus, human rotavir?, astrovir?, enterovir?, poliovir?, respiratory, hepatitis a, sabin